

บรรณานุกรม

บรรณานุกรม

- American Public Health Association. (1995). *Standard methods for the examination of water and wastewater* (19th ed). Washington, DC: American Public Health Association.
- American Society for Testing and Materials. (1997). *Annual book of ASTM standard: Water and environmental technology*. Philadelphia: American Society for Testing and Materials.
- AOAC International. (1993). *AOAC Peer-Verified Methods program*. Arlington: AOAC International.
- Bruijnsvoort, M. V., Sanghi, S. K., Poppe, H., & Kok, W. T. (1997). Determination of chloro phenols by micellar electrokinetic chromatography with electrochemical detection. *Journal of Chromatography A*, 757, 203-213.
- Chao, Y. C., & Whang, C. W. (1994). Capillary zone electrophoresis of eleven priority phenols with indirect fluorescence detection. *Journal of Chromatography A*, 663, 229-237.
- Chien, R., & Burgi, D. S. (1992). Sample stacking of an extremely large inject volume in high-performance capillary electrophoresis. *Analytical Chemistry*, 64, 1046-1050.
- Coutts, R. T., Hargesheimer, E. E., & Pasutto, F. M. (1979). Gas chromatographic analysis of trace phenols by direct acetylation in aqueous solution. *Journal of Chromatography*, 179, 291-299.
- Deng, Y. et al. (1998). Separation & determination of aromatic acids in natural water with preconcentration by capillary zone electrophoresis. *Journal of Chromatography A*, 817, 145-152.
- Fedorak, P. M., & Hrudey, S. E. (1988). Anaerobic degradation of phenolic compounds with application to treatment of industrial waste water. In D. L. Wise (Ed.), *Biotreatment systems* (pp. 169 – 226). Florida: CRC Press.
- Fung, Y. S. & Long, Y. H. (2001). Determination of phenols in soil by supercritical fluid extraction capillary electrochromatography. *Journal of Chromatography A*, 907, 301-311.
- Gaitonde, C. D., & Pathak, P. V. (1990). Capillary zone electrophoretic separation of chlorophenols in industrial waste water with on-column electrochemical detection *Journal of Chromatography*, 514, 389-393.

- Gawdzik, B., Gawdzik, J., & Czerwinska-Bil, U. (1990). Use of polymeric sorbent for the off-line preconcentration of priority pollutant phenols from water for high - performance liquid chromatography. *Journal of Chromatography*, 509, 135-140.
- Gonnord, M. F., & Collet, J. (1993). Optimized capillary zone electrophoretic separation of chlorinated phenols. *Journal of Chromatography*, 645, 327-336.
- Hilmi, A., Luong, J. H. T., & Ngnyen, A. L. (1997). Capillary electrophoretic separation of chlorophenols using amperometric detection. *Journal of Chromatography A*, 761, 259 - 268.
- Jandik, P., & Boon, G. (1993). *Capillary electrophoresis of small molecules and ions*. New York: VCH.
- Jorgenson, J. W., & Lukacs, K. D. (1981). Zone electrophoresis in open-tubular glass capillaries. *Analytical Chemistry*, 53, 1298 - 1302.
- Khan, K. A., Suidan, M. T., & Cross, W. H. (1981). Anaerobic activated carbon for the treatment of phenol-bearing wastewater. *Journal of Water Pollution Control Federation*, 53, 1519- 1532.
- Kumaran, P., & Shivaraman, N. (1988). Biological treatment of toxic industrial wastes. In D. L. Wise (Ed.), *Biotreatment systems* (pp. 227 - 284). Florida: CRC Press.
- Li, G., & Locke, D. C. (1995). Separation of the eleven priority pollutant phenols by capillary zone electrophoresis. *Journal of Chromatography B*, 669, 93-102.
- Lin, C. E., Lin, W. C., & Chion, W. C. (1995). Migration behavior & optimization of selectivity of dichlorophenols in capillary zone electrophoresis. *Journal of Chromatography A*, 705, 325-333.
- Makuch, B., Gazda, K., & Kaminski, M. (1993). Determination of phenol & monochlorophenols in water by reversed-phase liquid chromatography. *Analytical Chimica Acta*, 284, 53 - 58.
- Maman, O., Marseille, F., Guillet, B., Disnar, J. R., & Morin, P. (1996). Separation of phenolic aldehydes, ketones and acid from lignin degradation by capillary zone electrophoresis. *Journal of Chromatography A*, 755, 89-97.

- Mathew, J., & Elzerman, A. W. (1981). Gas-liquid chromatographic determination of some chloro and nitrophenols by direct acetylation in aqueous solution. *Analytical Letters*, 14(A16), 1351 – 1361.
- Ong, C. P., Ng, C. L., Chong, N. C., Lee, H. K., & Li, S. F. Y. (1990). Retention of eleven priority phenols using micellar electrokinetic chromatography. *Journal of Chromatography A*, 516, 263-270.
- Onodera, S., Nagatsuka, A., Rokuhara, T., Asakura, T., Naoko, H., & Suzuki, S. (1993). Re-evaluation of solid-phase adsorption and desorption techniques for isolation of trace organic pollutants from chlorinated water. *Journal of Chromatography*, 642, 185 – 194.
- Pocurull, E., Calull, M., Marce, R. M., & Burrull, F. (1996). Determination of phenolic compounds at low $\mu\text{g l}^{-1}$ levels by various solid-phase extractions followed by liquid chromatography and diode-array detection. *Journal of Chromatography A*, 719, 105 – 112.
- Rodriguez, I., Turnes, M. I., Bollain, M. H., Mejuto, M. C., & Cela, R. (1997). Determination of phenolic pollutants in drinking water by capillary electrophoresis in the sample stacking mode. *Journal of Chromatography A*, 778, 279–288.
- Rose, D. J., Jr., & Jorgenson, J. W. (1988). Characterization and automation of sample introduction method for capillary zone electrophoresis. *Analytical Chemistry*, 60(7), 642- 648.
- Skoog, D. A., West, D. M., & Holler, F. J. (1991). Fundamentals of analytical chemistry. Fort Wort: Horcourt Brace College.
- Song, X., & Budde, W. L. (1998). Determination of chlorinated acid herbicides and related compounds in water by capillary electrophoresis electrospray negative ion mass spectrometry. *Journal of Chromatography A*, 829, 327–340.
- Turnes, M. I., Mejuto, M. C., & Cela, R. (1996). Determination of pentachlorophenol in water samples by capillary zone electrophoresis. *Journal of Chromatography A*, 733, 395–404.
- Weinberger, R. (1993). *Practical capillary electrophoresis*. Boston: Academic Press.
- Zar, J. H. (1984). Biostatistical analysis (2nd ed). New Jersey: Prentice-Hill International, Inc.