

บรรณานุกรม

- ถัดดา แก้วกล้าปัญญาเจริญ (2544). การวิเคราะห์ปริมาณสารชาลบูตามอลตกค้างในเนื้อ ดับ และ ไตสุกร โดยวิธี HPLC. วารสารกรมวิทยาศาสตร์การแพทย์, 3, 215-221.
- ลีณา ลีลาพัฒน์กิจ (2542). การประยุกต์แคปิลารีอิเล็กโทรโฟรีซิสสำหรับการวิเคราะห์อาหาร. วารสารกรมวิทยาศาสตร์การแพทย์, 41(2), 221-229.
- สมทรง เลขะกุล (2542). ชีวเคมีของวิตามิน (พิมพ์ครั้งที่ 1). กรุงเทพฯ: ศกวนิชการพิมพ์.
- Ackermans, M. T., Beckers, J. L., Everaerts, F. M., & Seelen, I. G. J. A. (1992). Comparison of isotachopheresis, capillary zone electrophoresis and high-performance liquid chromatography for the determination of salbutamol, terbutaline sulphate and fenoterol hydrobromide in pharmaceutical dosage forms. *Journal of Chromatography*, 590, 341-353.
- Altria, K. D. (1993). Determination of salbutamol-related impurities by capillary electrophoresis. *Journal of Chromatography*, 634, 323-328.
- Andrés-Lacueva, C., Mattivi, F., & Tonon, D. (1998). Determination of riboflavin, flavin mononucleotide and flavin adenine dinucleotide in wine and other beverages by high-performance liquid chromatography with fluorescence detection.. *Journal of Chromatography A*, 823, 355-363.
- Barna, E., & Dworschák, E. (1994). Determination of thiamine (vitamin B₁) and riboflavin (vitamin B₂) in meat and liver by high-performance liquid chromatography. *Journal of Chromatography A*, 668, 359-363.
- Bilic, N., & Sieber, R. (1990). Determination of flavins in dairy products by high-performance liquid chromatography using sorboflavin as internal standard. *Journal of Chromatography*, 511, 359-366.
- Blomgren, A., & et al. (2002). Extraction of clenbuterol from calf urine using a molecularly imprinted polymer followed by quantitation by high-performance liquid chromatography with UV detection. *Journal of Chromatography A*, 975, 157-164.
- Camilleri, P. (1993). *Capillary electrophoresis: Theory and practice*. Florida: CRC Press.

- Capo-chichi, C. D., & et al. (2000). Analysis of riboflavin and riboflavin cofactor levels in plasma by high-performance liquid chromatography. *Journal of Chromatography B*, 739, 219-224.
- Carducci, C. N., Lucangioli, S. E., Rodríguez, V.G., & Otero, G.C.F. (1996). Application of extraction disks in dissolution tests of clenbuterol and levothyroxine tablets by capillary electrophoresis. *Journal of Chromatography A*, 730, 313-319.
- Cataldi, T. R. I., Nardiello, D., Benedetto, G. E. D., & Bufo, S. A. (2002). Optimizing separation conditions for riboflavin, flavin mononucleotide and flavin adenine dinucleotide in capillary zone electrophoresis with laser-induced fluorescence detection. *Journal of Chromatography A*, 968, 229-239.
- Cataldi, T. R. I., Nardiello, D., Scrano, L., & Scopa, A. (2002). Assay of riboflavin in sample wines by capillary zone electrophoresis and laser-induced fluorescence detection. *Journal of Agricultural and Food Chemistry*, 50, 6643-6647.
- Esquisable, A., & et al. (1997). Determination of salbutamol enantiomers by high-performance capillary electrophoresis and its application to dissolution assays. *Journal of Pharmaceutical and Biomedical Analysis*, 16, 357-366.
- Fiori, M., Cartoni, C., Bocca, B., & Brambilla, G. (2002). The use of nonendcapped C18 columns in the cleanup of clenbuterol and a new adrenergic agonist from bovine liver by gas chromatography-tandem mass spectrometry analysis. *Journal of Chromatographic Science*, 40, 92-96.
- Gausepohl, C., & Blaschke, G. (1998). Stereoselective determination of clenbuterol in human urine by capillary electrophoresis. *Journal of Chromatography B*, 713, 443-446.
- Gliszczynska, A., & Koziolowa, A. (1998). Chromatographic determination of flavin derivatives in baker's yeast. *Journal of Chromatography A*, 822, 59-66.
- Gliszczynska-Swinglo, A., & Koziolowa, A. (2000). Chromatographic determination of riboflavin and its derivatives in food. *Journal of Chromatography A*, 881, 285-297.
- González, P., & et al. (1997). Determination of residues of the β -agonist clenbuterol in liver of medicated farm animals by gas chromatography-mass spectrometry using diphasic dialysis as an extraction procedure. *Journal of Chromatography B*, 693, 321-326.

- Grossman, P. D., & Colburn, J. C. (1992). *Capillary electrophoresis. theory and practice*. San Diego: Eds., Academic Press.
- Huber, L. (1998). *Validation and qualification in analytical laboratories*. Buffalo Grove, IL: Interpharm Press.
- Jandik, P., & Bonn, G. (1993). *Capillary electrophoresis of small molecules and ions*. New York: VCH.
- Koh, Y. M., Saleh, M. I., & Tan, S. C. (2003). Selective extraction of salbutamol from human plasma with the use of phenylboronic acid. *Journal of Chromatography A*, 987, 257-267.
- Koole, A., Bosman, J., Franke, J. P., & Zeeuw, R. A. D. (1999). Multiresidue analysis of β_2 -agonists in human and calf urine using multimodal solid-phase extraction and high-performance liquid chromatography with electrochemical detection. *Journal of Chromatography B*, 726, 149-156.
- Lawrence, J. F., & Ménard, C. (1997). Determination of clenbuterol in beef liver and muscle tissue using immunoaffinity chromatographic cleanup and liquid chromatography with ultraviolet absorbance detection. *Journal of Chromatography B*, 696, 291-297.
- Mälkki, L., & Tammilehto, S. (1990). Decomposition of salbutamol in aqueous solutions. I. The effect of pH, temperature and drug concentration. *International Journal of Pharmaceutics*, 63, 17-22.
- Mälkki-Laine, L., & Hartikainen, E. (1996). Electrokinetic behaviour of salbutamol and its decomposition products and determination of salbutamol by micellar electrokinetic capillary chromatography. *Journal of Chromatography A*, 724, 297-306.
- Mälkki-Laine, L., Purra, K., Kähkönen, K., & Tammilehto, S. (1995). Decomposition of salbutamol in aqueous solutions. II. The effect of buffer species, pH, buffer concentration and antioxidants. *International Journal of Pharmaceutics*, 117, 189-195.
- Mattivi, F., Monetti, A., Vrhovšek, U., Tonon, D., & Andrés-Lacueva, C. (2000). High-performance liquid chromatographic determination of the riboflavin concentration in white wines for predicting their resistance to light. *Journal of Chromatography A*, 888, 121-127.

- Moreno, P., & Salvadó, V. (2000). Determination of eight water- and fat-soluble vitamins in multi-vitamin pharmaceutical formulations by high-performance liquid chromatography. *Journal of Chromatography A*, 820, 207-215.
- Natakankitkul, S. (1997). Separation of clenbuterol enantiomers by cyclodextrin-modified capillary electrophoresis. *Thai Journal Pharmaceutical Science*, 21, 191-198.
- Posyniak, A., Zmudzki, J., & Niedzielska, J. (2003). Screening procedures for clenbuterol residue determination in bovine urine and liver matrices using enzyme-linked immunosorbent assay and liquid chromatography. *Analytica Chimica Acta*, 483, 61-67.
- Ramos, F. J. D. (2000). β_2 -Agonist extraction procedures for chromatographic analysis. *Journal of Chromatography A*, 880, 2000, 69-83.
- Russell, L.F., Brooks, L., & McRae, K.B. (1998). Development of a robotic-HPLC determination of riboflavin vitamers in food. *Food Chemistry*, 63, 125-131.
- Shintani, H., & Polonský, J. (1997). *Handbook of electrophoresis applications*. London: Chapman & Hall. 18.
- Song, Y., & et. al. (2003). Direct separation and quantitative determination of clenbuterol enantiomers by high performance liquid chromatography using an amide type chiral stationary phase. *Journal of Pharmaceutical and Biomedical Analysis*, 31, 311-319.
- Srichana, T., & Suedee, R. (2000). Enantiomeric separation of salbutamol from spiked urine extracted with solid stationary phase. *Thai Journal Pharmaceutical Science*, 24 (1), 27-36.
- Stol, R., Mazereeuw, M., Tjaden, U. R., & Greef, J. V. D. (2000). Pseudo-electrokinetic packing of high efficiency columns for capillary electrochromatography. *Journal of Chromatography A*, 873, 293-298.
- Su, A.-K., & Lin, C.-H. (2003). Determination of riboflavin in urine by capillary electrophoresis-blue light emitting diode-induced fluorescence detection combined with a stacking technique. *Journal of Chromatography B*, 785, 39-46.
- Traynor, I. M., Crooks, S. R. H., Bowers, J., & Elliott, C. T. (2003). Detection of multi- β -agonist residues in liver matrix by use of a surface plasma resonance biosensor. *Analytica Chimica Acta*, 483, 187-191.

Weinberger, R. (1993). *Practical capillary electrophoresis*. New York: Harcourt Brace Jovanovich. 19-200.

Zhou, T., Hu, Q., Yu, H., & Fang, Y. (2001). Separation and determination of β -agonists in serum by capillary zone electrophoresis with amperometric detection. *Analytica Chimica Acta*, 441, 23-28.

มหาวิทยาลัยบูรพา
Burapha University