

บรรณานุกรม

- กองโภชนาการ กรมอนามัย กระทรวงสาธารณสุข. (2546). รายงานการสำรวจภาวะอาหารและโภชนาการของประเทศไทย ครั้งที่ 5. กรุงเทพฯ: กองโภชนาการ.
- จันทนา เชื้อนาค. (2542). การศึกษาเปรียบเทียบการเปลี่ยนแปลงของระดับไขมันในเลือดและน้ำหนักตัวในผู้สูงอายุเมื่อใช้และไม่ใช้คู่มือ การเฝ้าระวังการบริโภคอาหารไขมัน. *เวชสาร.*, 43(2), 135-141.
- เจลิยว ปิยะชน. (2547). หลอดเลือดแข็งตีตัน หัวใจ, สมอ, ... อื่น ๆ: พื้นดินสภาพด้วยการแพทย์องค์รวม. กรุงเทพฯ: สุขภาพใจ
- นิธิยา รัตนปนนท์. (2549). เคมีอาหาร (พิมพ์ครั้งที่ 2). กรุงเทพฯ: โอเคียนสโตร์
- พรทิพย์ โล่เลขา. (2536). ไลโปโปรตีนและภาวะหลอดเลือดแดงแข็ง. กรุงเทพฯ: ชัยเจริญ.
- ภัทรารุช อินทรกำแหง. (ม.ป.ป.). *Exercise and dyslipidemia*. ม.ป.ท.
- วรรณภา วัฒนกุล. (2541). ปริมาณไขมันในเลือดของนักเดินแอโรบิกหญิง. วิทยานิพนธ์ปริญญาวิทยาศาสตรมหาบัณฑิต, สาขาวิทยาศาสตร์การออกกำลังกายและการกีฬา, บัณฑิตวิทยาลัย, มหาวิทยาลัยบูรพา.
- Aellen, R., Hollmann, W., & Boutellier, U. (1993). Effects of aerobic and anaerobic training on plasma lipoproteins, *J. Sports Med.*, 14, 396-400.
- American College of Sports Medicine. (1998). The recommended Quantity and quality for exercise for developing and maintaining cardiorespiratory and muscular fitness in healthy adults. *Med. Sc.i Sports Exere.*, 30(6), 265-74.
- Aquilo, A., Tauler, P., Guix, M. P., Villa, G., Cordova, A., Tur, J. A., & Pons, A. (2003). Effect of exercise intensity and training on antioxidants and cholesterol profile in cyclists. *Journal of Nutrition Biochemistry*, 14, 319-325.
- Barter, P. (2004). Is high-density lipoprotein the protector of the cardiovascular system? *European Heart Journal Supplements*, 6 (supplement A), A19-A22
- Belsey, J. (1998). Lipid-lowering in coronary heart disease. *Gavel*, 1(1), 1-8.
- Boutcher, S. H., Meyer, B. J., Craig, G. A., & Astheimer, L. (2003). Plasma lipid and fibrinogen levels in aerobically trained and untrained postmenopausal women. *J Sports Med Phys Fitness*, 43, 231-235.

- Brooks, G. A., Fahey, T. D., White, T. P., & Baldwin, K. M. (1999). *Exercise physiology: Human bioenergetics and its applications* (3rd ed.). New York: McGraw-Hill.
- Brownell, K. D., Bachorik, P. S., & Ayerle, R. S. (1982). Changes in plasma lipid and lipoprotein levels in men and women after a program of moderate exercise. *Circulation*, 65(3), 477-484.
- Christ, M., Iannello, C., Iannello, P., & Grimm, W. (2004). Effects of a weight reducing program with and without aerobic exercise in the metabolic syndrome. *International Journal of Cardiology*, 97, 115-122.
- Christie, W. W. (1993). In *Advances in Lipid Methodology* (2nd ed.). Dundee: W.W. Christie, Oily Press.
- Crouse, S. F., O'Brien, B. C., Grandjean, P. W., Lowe, R. C., Rohack, J. J., Green, J. S., & Tolson, H. (1997). Training intensity, blood lipids, and apolipoproteins in men with high cholesterol. *J. Appl. Physiol.*, 82(1), 270-277.
- Crouse, S. F., O'Brien, B. C., Rohack, J. J., Lower, R. C., Green, J. S., Tolson, H., & Reed, J. L. (1995). Changes in serum lipids and apolipoproteins after exercise in men with high cholesterol: influence of intensity. *J. Appl. Physiol.*, 79(1), 279-286.
- Davis, P. S., Bartoli, W. P., & Dustine, J. L. (1993). Effect of acute exercise intensity on plasma lipids and apolipoproteins in trained runners. In *Yearbook of Sports Medicine 1993*. St. Louis, M. O.: Mosby-Year Book.
- Dowling, E. A. (2001). How exercise affects lipid profiles in women: what to recommend for patients. *The Physician and Sportsmedicine*, 29(9), 45-52.
- Durstine, J. L., & Haskell, W. L. (1994). Effects of exercise training on plasma lipids and lipoproteins. *Exercise and Sports Science Reviews*, 22, 477-522.
- Ferrauti, A., Weber, K., & Struder, H. K. (1997). Effects of tennis training on lipid metabolism and lipoproteins in recreational players. *Br. J. Sport Med.*, 31, 322-327.
- Forge, R. (2001). Managing cholesterol with exercise. ACE Fit Facts. Retrieved March 7, 2007, from <http://www.acefitness.org>
- Friedewald, W. T., Levy, R. I., & Fredrickson, D. S. (1972). Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin. Chem.*, 18, 499-502.

- Gordon, P. M., Fowler, S., Warty, V., Danduran, M., Visich, P., & Keteyian, S. (1998). Effects of acute exercise on high density lipoprotein cholesterol and high density lipoprotein subfractions in moderately trained females. *Br. J. Sports Med.*, 32, 63-67.
- Hainline, A. Jr., Karon, J., & Lippel, K. (Eds.). (1982). *Manual of laboratory operations: lipid and lipoprotein analysis* (2nd ed). Rev. Washington, D.C.: Government printing Office. (DHEW publication no. (NIH) 75-628.)
- Hardman, A. E. (1998). The influence of exercise on postprandial triacylglycerol metabolism. *Atherosclerosis*, 141, S93-S100.
- Haswell, W. L. (1984). The influence of exercise on the concentration of triglyceride and cholesterol in human plasma. *Exer. Sport. Sci. Rev.*, 12, 205-24.
- Hicks, A. L., MacDougall, J. D., & Muckle, T. L. (1987). Acute changes in high-density lipoprotein cholesterol with exercise of different intensities. *J. Appl. Physiol.*, 63(5), 1956-1960.
- Horowitz, J. F., & Klein, S. (2000). Lipid metabolism during endurance exercise. *Am J Clin Nutr*, 72(suppl), 558S-563S.
- Horowitz, J. F. (2003). Fatty acid mobilization from adipose tissue during exercise. *TRENDS in Endocrinology and Metabolism*, 14(8), 386-392.
- Isler, A. K., Kosar, S. N., & Korkusuz, F. (2001). Effects of step aerobics and aerobic dancing on serum lipids and lipoproteins. *J. Sport Med. Phys. Fitness*, 41, 380-385.
- Kantor, M. A., Cullinane, E. M., Sady, S. P., Herbert, P. N., & Thompson, P. D. (1987). Exercise acutely increases high-density lipoprotein-cholesterol and lipoprotein lipase activity in trained and untrained men. *Metabolism*, 36(2), 188-192.
- Katzmarzyk, P. T., Leon, A. S., Rankinen, T., Gagnon, J., Skinner, J. S., Wilmore, J. H., Rao, D. C., & Bouchard, C. (2001). Changes in blood lipids consequent to aerobic exercise training related to changes in body fatness and aerobic fitness. *Metabolism*, 50(7), 841-848.
- Kravitz, L., & Heyward, V. (1994). The exercise & cholesterol controversy. *IDEA today*, 12(2), 38-42.

- Lakka, H., Tremblay, A., Despres, J., & Bouchard, C. (2004). Effects of long-term negative energy balance with exercise on plasma lipid and lipoprotein levels in identical twins. *Atherosclerosis*, 172, 127-133.
- Madsen, P. (2004). Exercise and high-density lipoprotein: the effects on coronary heart disease risk. *TSMJ*, 5, 11-16.
- Manore, M., & Thompson, J. (2000) *Sport nutrition for health and performance*. Champaign, IL: Human kinetics.
- Marshall, W. J. (1992). *Illustrated textbook of clinical chemistry* (2nd ed.). London: Gowre Medical Pub.
- Marti, B., Suter, E., Riesen, W. F., Tschopp, A., Wanner, H. U., & Gutzwiller, F. (1990). Effects of long-term, self-monitored exercise on the serum lipoprotein and apolipoprotein profile in middle-aged men. *Atherosclerosis*, 81, 19-31.
- Mougiou, V., Ring, S., Petridou, A., & Nikolaidis, M. (2003). Duration of coffee- and exercise-induced changes in the fatty acid profile of human serum. *J Appl Physiol*, 94, 476-484.
- Neibauer, J., Hambrecht, R., Velich, T., Marburger, C., Hauer, K., Kreuzer, J., Zimmermann, R., Hadenberg, E., Schlierf, G., Schuler, G., & Kubler, W. (1996). Predictive value of lipid profile for salutary coronary angiographic changes in patients on a low-fat diet and physical exercise program. *The American Journal of Cardiology*, 78, 163-167.
- Nelson, D. L. & Cox, M. M. (2000). *Lehninger principles of biochemistry* (3rd ed.). New York: Worth.
- Nicklas, B. J., Katzel, L. I., Busby-Whitehead, J., & Goldberg, A. P. (1997). Increases in high-density lipoprotein cholesterol with endurance exercise training are blunted in obese compared with lean men. *Metabolism*, 46(5), 556-561.
- Nobel, J. B. (1986). *Physiology of exercise and sport*. St. Louis: Times Mirror/Mosby.
- Powers, S. K., & Howley, E. T. (2001). *Exercise Physiology: theory and application to fitness and performance* (4th ed.). Boston: McGraw-Hill.
- Rajman, J., Kendall, M. J., Holder, R. L., Salih, M., & Gammage, M. D. (1996). Investigation of low density lipoprotein subfractions as a coronary risk factor in normotriglyceridaemic men. *Atherosclerosis*, 125, 231-242.

- Sanchez-Quesada, J. L., Ortega, H., Payes-Romero, A., Serrat-Serrat, J., Gonzalez-Sastre, F., Lasuncion, M. A., & Ordonez-Llanos, J. (1997). LDL from aerobically-trained subjects shows higher resistance to oxidative modification than LDL from sedentary subjects. *Atherosclerosis*, 132, 207-213.
- Sgouraki, E., Tsopanakis, A., & Tsopanakis, C. (2001). Acute exercise: Response of HDL-C, LDL-C lipoproteins and HDL-C subfractions levels in selected sport disciplines. *J. Sports Med. Phys. Fitness*, 41, 386-391.
- Stefanick, M. L., Mackey, S., Sheehan, M., Ellsworth, N., Haskell, W., & Wood, P. D. (1998). Effects of diet and exercise in men and postmenopausal women with low levels of HDL cholesterol and high levels of LDL cholesterol. *The New England Journal of Medicine*, 339(1), 12-20.
- Stein, R. A., Michielli, D. W., Glantz, M. D., Sardy, H., Cohen, A., Goldberg, N., & Brown, C. D. (1990). Effect of different exercise Training intensities on lipoprotein cholesterol fractions in healthy middle aged men. *Am. Heart. J.*, 119, 277-283.
- Sunami, Y., Motoyama, M., Kinoshita, F., Mizooka, Y., Sueta, K., Matsunaga, A., Sasaki, J., Tataka, H., & Shindo, M. (1999). Effects of low-intensity aerobic training on the high-density lipoprotein cholesterol concentration in healthy elderly subjects. *Metabolism*, 48(8), 984-988.
- Thaikruea, L., Seetarnanotch, W., & Seetarnanotch, S. (2006). Appropriate cut-off level of BMI for screening in Thai adults. *J Med Assoc Thai.*, 89(12), 2123-2128.
- Thompson, P. D., Cullinane, E. M., Dady, S. P., Flynn, M. M., Bernier, D. N., Kantor, M. A., Saritelli, A. L., & Herbert, P. N. (1988). Modest changes in high-density lipoprotein concentration and metabolism with prolonged exercise training. *Circulation*, 78(1), 25-34.
- Thompson, P. D., Yurgalevitch, S. M., Flynn, M. M., Zmuda, J. M., Spannaus-Martin, D., Saritelli, A., Bausserman, L., & Herbert, P. N. (1997). Effect of prolonged exercise training without weight loss on high-density lipoprotein metabolism in overweight men. *Metabolism*, 46(2), 217-223.

- Tsai, A. C., Sandretto, A., & Chung, Y. (2003). Dieting is more effective in reducing weight but exercise is more effective in reducing fat during the early phase of a weight-reducing program in healthy humans. *Journal of Nutritional Biochemistry*, 14, 541-549.
- Tsai, J., Liu, J., Kao, C., Tomlinson, B., Kao, P., Chen, J., & Chan, P. (2002). Beneficial effects on blood pressure and lipid profile of programmed exercise training in subjects with white coat hypertension. *AJH*, 15, 571-576.
- Vella, C. A., Kravitz, L., & Janot, J. M. (2001). A review of the impact of exercise on cholesterol levels. *IDEA Health & Fitness Source*, 19(10), 48-54.
- Warnick, G. R., Benderson, J., & Albers, J. J. (1982). Dextran sulfate-Mg²⁺ precipitation procedure for quantitation of high-density-lipoprotein cholesterol. *Clin. Chem.*, 28, 1379-1388.
- White, D. A., Middleton, B., & Baxter, M. (1984). *Hormone and metabolic control*. London: Edward Arnold.