

บรรณานุกรม

- กรมควบคุมมลพิษ และสมาคมวิศวกรสิ่งแวดล้อมแห่งประเทศไทย. (2537). *รายละเอียดสนับสนุนเกณฑ์แนะนำการออกแบบระบบรวบรวมน้ำเสียและโรงปรับปรุงคุณภาพน้ำของชุมชน*. กรุงเทพฯ: โรงพิมพ์เรือนแก้วการพิมพ์.
- กรมโรงงานอุตสาหกรรม. (2545). *ตำราระบบบำบัดมลพิษน้ำ*. กรุงเทพฯ: สมาคมวิศวกรรมสิ่งแวดล้อมแห่งประเทศไทย.
- เกรียงศักดิ์ อุคมสินโรจน์. (2539). *การบำบัดน้ำเสีย*. กรุงเทพฯ: มิตรนราการพิมพ์.
- เกรียงศักดิ์ อุคมสินโรจน์. (2543). *วิศวกรรมการกำจัดน้ำเสียเล่ม 4*. กรุงเทพฯ: มิตรนราการพิมพ์.
- เกรียงศักดิ์ อุคมสินโรจน์. (2550). *วิศวกรรมสิ่งแวดล้อม*. กรุงเทพฯ: สยามสเตชันเนอรีซ์พลาซัส.
- ณฐนนท์ ตราชู. (2551). *จุลินทรีย์ในแผ่นชีวะ*. มหาสารคาม: อภิชาติการพิมพ์.
- คุณภา คำเพชร. (2548). *การบำบัดน้ำเสียด้วยระบบบำบัดน้ำเสียแบบจุลินทรีย์แขวนลอยและจุลินทรีย์บนตัวกลางยึดเกาะ*. วิทยานิพนธ์ปริญญาวิทยาศาสตรมหาบัณฑิต, สาขาวิชาวิทยาศาสตร์สิ่งแวดล้อม, คณะวิทยาศาสตร์, มหาวิทยาลัยบูรพา.
- ธงชัย พรรณสวัสดิ์. (2530). *น้ำเสียชุมชนและปัญหามลภาวะทางน้ำในเขต กทม. และปริมณฑล*. กรุงเทพฯ : โรงพิมพ์เรือนแก้วการพิมพ์.
- นิชุตนา บินนิมะ. (2549). *การบำบัดน้ำเสียด้วยจุลินทรีย์แบบยึดเกาะตัวกลางชนิดเส้นใยในถ่อ* โดยระบบเลี้ยงตะกอน. วิทยานิพนธ์ปริญญาวิศวกรรมศาสตรมหาบัณฑิต, สาขาวิชาวิศวกรรมสิ่งแวดล้อม, คณะวิศวกรรมศาสตร์, มหาวิทยาลัยเกษตรศาสตร์.
- บัญญัติ สุขศรีงาม. (2534). *จุลชีววิทยาทั่วไป*. กรุงเทพฯ: สำนักพิมพ์โอเดียนสโตร์.
- Alleman, J. E. (1984). Elevated nitrite occurrence in biological wastewater treatment systems. *Wat. Sci. Technol.*, 17, 409-419.
- Antoniou, P., Hamilton, J., Koopman, B., Jain, R., Holloway, B., Lyberatos, G., & Svoronos, S. A. (1990). Effect of temperature and pH on the effective maximum specific growth rate of nitrifying bacteria. *Wat. Res.*, 24, 97-101.
- Bejar, V., Llamas, I., Calvo, C., & Quesada, E. (1998). Characterization of exopolysaccharides produced by 19 halophilic strains of the species *Halomanas eurihalina*. *J. Biotech.*, 61, 135-141.
- Brown, J. H., & Lester, J.N. (1980). Comparison of bacterial extracellular polymer extraction methods. *Appl. Envir. Microbiol.*, 40, 179-185.

- Bura, R., Cheng, M., Liao, B., Finlayson, J., Lee, B.C., Droppo, I.G., Leppard, G.G., & Liss, S.N. (1998). Composition of extracellular polymeric substances in the activated sludge floc matrix. *Wat. Sci. Tech.*, 37(4-5), 325-333.
- Busch, P., & Stumm, W. (1968). Chemical interactions in the aggregation of bacteria, bioflocculation in waste treatment. *Environ. Sci. Technol.*, 2(1), 49-53.
- Chao, A.C., & Keinath, T.M. (1979). Influence of process loading intensity on sludge clarification and thickening characteristics. *Wat. Res.*, 13(12), 1213-1220.
- Dunne, W.M., & Burd, E.M. (1992). The Effects of magnesium, calcium, EDTA, and pH on the In vitro adhesion of staphylococcus epidermidis to plastic. *Microbiol Immunol.* 1992,36(10),1019-1027.
- Durmaz, B., & Sanin, F.D. (2001). Effect of carbon to nitrogen ratio on the composition of microbial extracellular polymers in activated sludge. *Wat. Sci. and Technol.*, 44(10), 221-229.
- Durmaz, B., & Sanin, F.D. (2001). Effect of carbon to nitrogen ratio on the physical and chemical properties of activated sludge. *Env.Tech.*, 24, 1331-1340.
- Friedman, B. A., Dugan, P.R., Pfister, R.M., & Remsen, C.C. (1996). Structure of exocellular polymers and their relationship to bacterial flocculation. *J. Bacteriol.*, 98(3), 1328-1334.
- Fronlund, B., Palmgren, R., Kieding, K., & Nielsen, P.H. (1996). Extraction of extracellular polymers from activated sludge using cation exchange resin. *Water Res.*, 30, 1749-1758.
- Gehr, R., & Henry, J.G. (1983). Removal of extracellular materials, techniques and pitfalls. *Wat. Res.*, 17(12), 1743-1748.
- Gerardi, M.H. (2002). *Nitrification and denitrification in the activated sludge process*. Published simultaneously in Canada. United States of America.
- Giaouris, E., Chorianopoulos, N., & Nychas, G.-J.E. (2005). Effect of temperature, pH, and water activity on biofilm formation by *Salmonella enteric enteritidis* PT4 on stainless steel surfaces as indicated by the bead vortexing method and conductance measurements. *Journal of food protection*, 68, 2149-2154.

- Goodwin, J.A., & Foster, C.F. (1985). A further examination into the composition of activated sludge surfaces in relation to their settlement characteristics. *Wat. Res.*, 19, 527-533.
- Gulas, V., Bond, M., & Benefield, L. (1979). Use of exocellular polymers for thickening and dewatering activated sludge. *J. Water Pollut. Control Fed.*, 51(4), 798-807.
- Higgins, M.J., & Novak, J.T. (1997a). The effect of cation on settling and dewatering of activated sludge: Laboratory results. *Wat. Environ. Res.*, 69, 215-224.
- Higgins, M.J., & Novak, J.T. (1997b). Dewatering and settling of activated sludge: the case for using cation analysis. *Wat. Environ. Res.*, 69, 225-232.
- Higgins, M.J., & Novak, J.T. (1997c). Characterization of exocellular protein and its role in bioflocculation. *J. Env. Eng.*, 123(5), 479-485.
- Hoa, P.T., Nair, L., & Visvanathan, C. (2003). The effect of nutrients on extracellular polymeric substance production and its influence on sludge properties. *Wat. SA.*, 29(4), 437-442.
- Horan, N.J., & Eccles, C.R. (1986). Purification and characterization of extracellular polysaccharide from activated sludges. *Wat. Res.*, 20, 1427-1432.
- Janette Tournay & Bryne T. Ngwenya. (2010). The effect of ionic strength on the electrophoretic mobility and protonation constants of an EPS-producing bacterial strain. *Journal of Colloid and Interface Scienc.*, 348, 348-354.
- Jang, N., Ren, X.H., Kim, G., Ahn, C.H., Cho, J., & Kim, I.S. (2007). Characteristics of soluble microbial products and extracellular polymeric substances in the membrane bioreactor for water reuse. *Desalination*, 202(1-3), 90-98.
- Johnson, T., Steichen, M., Shaw, A., McQuarrie, J., & Hunter, G. (2006). When is IFAS the right choice. *Wat. Environ Foundation*, 200-224.
- Jones, G. L., & Paskins, A. R. (1982). Influence of high partial pressure of carbon dioxide and/or oxygen on nitrification. *J. Chem. Technol. Biotechnol.*, 32, 213-223.
- Jorand, F., Boue-Bigne, F., Block, J. C., & Urbain, V. (1998). Hydrophobic/hydrophilic properties of activated sludge exopolymeric substances. *Wat. Sci. Tech.*, 37 (4-5), 307-315.
- Jorand, F., Zartarin, F., Thomas, F., Bloc, J.C., Bottero, J.Y., Villemin, G., Urbain, V., & Manem, J. (1995). Chemical and structure (2D) linkage between bacteria within activated sludge flocs. *Wat. Res.*, 29, 1639-1647.

- Kampfer, P., & Wagner, M. (2002). Filamentous bacteria in activated sludge: Current taxonomic status and ecology. *Encyclopedia of environmental microbiology*, 1287-1306.
- Laspidou, C.S., & Rittman, B.E. (2002). A unified theory for exopolymeric substances, soluble microbial products, and active and inert biomass. *Wat. Res.*, 36, 2711-2720.
- Lee, K.C., & Rittmann, B.E. (2003). Effect of pH and precipitation on autohydrogenotrophic denitrification using the hollow-fiber membrane-biofilm reactor. *Wat. Res.*, 37, 1551-1556.
- Liao, B.Q., Allen, D.G., Droppo, I.G., Leppard, G.G., & Liss, S.N. (2001). Surface properties of sludge and their role in bioflocculation and settleability. *Wat. Res.*, 35, 339-350.
- Li, D.H., & Ganczarczyk, J.J. (1990). Structure of activated sludge flocs. *Biotechnol. Bioeng.*, 35(1), 57-65.
- Ling-Ling Wang, Long-Fei Wang, Xue-Mei Ren, Xiao-Dong Ye, Wen-Wei Li, Shi-Jie Yuan Min Sun, Guo-Ping Sheng, Han-Qing Yu, & Xiang-Ke Wang. (2011). pH Dependence of Structure and Surface Properties of Microbial EPS. *Environ. Sci. Technol.*, 46, 737-744.
- Mckinny, R.E. (1952). *Biological treatment of sewage and industrial wastews*. Reinhold, NY, USA.
- Morgan, J.W., Forster, C.F., & Evisom, L. (1990). A Comparative study of the nature of biopolymers extracted from anaerobic and activated sludges. *Wat. Res.*, 24(6), 743-750.
- Nielsen, P.H., Jahn, A., & Palmgren, R. (1997). Conceptual model for production and composition of exopolymers in biofilms. *Wat. Sci. Tech.*, 44, 1231-1237.
- Painter, H.A., & Loveless, J.E. (1983). Effect of temperature and pH value on the growth-rate constants of nitrifying bacteria in the activated-sludge process. *Wat. Res.*, 17, 237-248.
- Pavoni, J.L., Tenney, M.W., & Echelberger, W.F. (1972). Bacterial exocellular polymers and biological phosphorus and biological flocculation. *J. Water Pollut. Control Fed.*, 44, 414-431.
- Yongzhen Peng & Guibing Zhu. (2006). Biological nitrogen removal with nitrification and denitrification via nitrite pathway. *Appl Microbiol Biotechnol.*, 73, 15-26.

- Ramli, N.S.K., Guan, C.E., Nathan, S., & Vadivelu, J. (2012). The effect of environmental condition on biofilm formation of *Burkholderia pseudomallei* clinical isolates. *PloS ONE*, 7(9), 1-10.
- Rittmann, B.E., & McCarty, P.L. (2001). *Environmental biotechnology: principles and applications*. Singapore: McGraw-Hill.
- Ruiz, G., Jeison, D., & Chamy, R. (2003). Nitrification with high nitrite accumulation for the treatment of wastewater with high ammonia concentration. *Wat. Res.*, 37, 1371-1377.
- Sesay, M.L., & Sanin, F.D. (2004). An investigation of activated sludge floc structure in relation to solids retention time. *J. Residuals Sci. Tech.*, 1(2), 125-131.
- Shammas, N.K. (1986). Interactions of temperature, pH, and biomass on the nitrification process. *J. Water Pollution Control Federation*, 58(1), 52-59.
- Sheintuch, M., Lev, O., Einav, P., & Rubin, E. (1986). The role of exocellular polymer in the design of activated sludge. *Biotechnol. Bioeng.*, 28, 1564-1576.
- Sponza, D.T. (2003). Investigation of extracellular polymer substance (EPS) and Physicochemical properties of different activated sludge flocs under steady-state conditions. *Enzyme Microb. Tech.*, 32, 375-385.
- Strous, M., Kuenen, J.G., & Jetten, M.S.M. (1999). Key physiology of anaerobic ammonium oxidation. *Applied and environmental microbiology*, 65, 3248-3250.
- Tchobanoglous, G., Burton, F.L., & Stensel, H.D. (1991). *Wastewater engineering: treatment and reuse*. New York, USA: McGraw-Hill.
- Tenney, M.W., & Stumm, W. (1965). Chemical flocculation of microorganisms in biological waste treatment. *J. Water Pollut Control Fed.*, 37, 1370-1388.
- Tezuka, Y. (1969). Cation-dependent flocculation in flavobacterium species predominant in activated sludge. *Appl. Microbial.*, 17(2), 222-226.
- Urbain, V., Block, J.C., & Manem, J. (1993). Bioflocculation in activated sludge: an analytic approach. *Wat. Res.*, 27(5), 829-838.
- Wilén, B.M., Jin, B., & Lant, P. (2003b). Relationship between flocculation of activated sludge composition of extracellular polymeric substance. *Wat. Sci. Tech.*, 47(12), 95-103.
- Wingender, J., Neu, T.R., & Flemming, H.C. (1999). *Microbial Extracellular Polymeric Substances: Characterisation, Structure and Function*. Springer, Berlin.